

A DIRECT PROOF OF $\mathcal{Z}$ -STABILITY FOR AH ALGEBRAS OF BOUNDED TOPOLOGICAL DIMENSION

MARIUS DADARLAT, N. CHRISTOPHER PHILLIPS, AND ANDREW S. TOMS

ABSTRACT. We prove that a unital simple approximately homogeneous (AH) C^* -algebra with no dimension growth absorbs the Jiang-Su algebra tensorially without appealing to the classification theory of these algebras. Our main result continues to hold under the slightly weaker hypothesis of exponentially slow dimension growth.

1. INTRODUCTION

The property of absorbing the Jiang-Su algebra $\mathcal{Z}$ tensorially— $\mathcal{Z}$ -stability, briefly—is a powerful regularity property for separable amenable C^* -algebras. It is a necessary condition for the confirmation of G. A. Elliott’s K-theoretic rigidity conjecture, which predicts that Banach algebra K-theory and positive traces will form a complete invariant for simple separable amenable C^* -algebras. We refer the reader to [9] for an up-to-date account of $\mathcal{Z}$ -stability as it relates to Elliott’s conjecture.

The necessity of $\mathcal{Z}$ -stability for K-theoretic classification suggests a two-step approach to further positive classification results: first, establish broad classification theorems for $\mathcal{Z}$ -stable C^* -algebras; second, prove that natural examples of simple separable amenable C^* -algebras are $\mathcal{Z}$ -stable. Winter, in a series of papers, has made significant contributions to the first part of this program. For instance, he has shown that the C^* -algebras associated to minimal uniquely ergodic diffeomorphisms satisfy Elliott’s conjecture modulo $\mathcal{Z}$ -stability. But there has so far been no progress on the second part of the program. This is not to say that we do not have natural examples of $\mathcal{Z}$ -stable C^* -algebras. It is only that the $\mathcal{Z}$ -stability of these examples is typically a consequence of having proved directly that the said examples satisfy Elliott’s conjecture.

If we are to have any hope of carrying out the suggested two-step approach to Elliott’s conjecture, then we must understand why already classified C^* -algebras are $\mathcal{Z}$ -stable *without* appealing to the heavy machinery of classification. The purpose of this article is to give a direct-read “not passing through classification”—proof that unital simple approximately homogeneous (AH) C^* -algebras with no dimension growth are $\mathcal{Z}$ -stable. The result that these C^* -algebras satisfy Elliott’s conjecture, due to various combinations of Elliott, Gong, and Li, is one of the most difficult theorems in the classification theory for separable stably finite

Date: June 17, 2008.

2000 Mathematics Subject Classification. Primary 46L35, Secondary 46L80.

Key words and phrases. Jiang-Su algebra, $\mathcal{Z}$ -stability, AH algebra, dimension growth.

C*-algebras, and is therefore an appropriate starting point for understanding $\mathcal{Z}$ -stability (see [6], [7], and [10]).

Finally, let us mention that W. Winter has recently announced a proof of $\mathcal{Z}$ -stability for a class of simple C*-algebras which includes the unital simple AH algebras of no dimension growth, using techniques which differ substantially from ours. Our result, however, allows one to relax the no dimension growth condition to a slightly weaker notion of “exponentially slow dimension growth”, and so is not subsumed by Winter’s result.

Acknowledgements. This work was carried out during the Fields Institute Thematic Program on Operator Algebras in the fall of 2007. The authors are grateful to that institution for its support. The first named author was partially supported by NSF grant #DMS-0500693

2. PRELIMINARIES

2.1. Generalities. We use M_n to denote the C*-algebra of $n \times n$ matrices with entries in $\mathbb{C}$. Let F and H be subsets of a metric space X and let $\epsilon > 0$ be given. We write $F \subseteq_\epsilon H$ if for each $f \in F$ there is some $h \in H$ such that $\text{dist}(f, h) < \epsilon$. We write $F \approx_\epsilon H$ if there is a bijection $\eta : F \rightarrow H$ such that $\text{dist}(f, \eta(f)) < \epsilon$.

2.2. AH algebras. An approximately homogeneous C*-algebra (*AH algebra*) is the limit of an inductive sequence $(A_i, \phi_i)_{i=1}^\infty$, where each A_i has the following form:

$$A_i = \bigoplus_{l=1}^{n_i} p_{i,l} (C(X_{i,l}) \otimes \mathcal{K}) p_{i,l},$$

where n_i is a natural number, $X_{i,l}$ is a compact metric space, $\mathcal{K}$ denotes the C*-algebra of compact operators on a separable infinite-dimensional Hilbert space, and $p_{i,l}$ is a projection in $C(X_{i,l}) \otimes \mathcal{K}$. C*-algebras of this form are called *semi-homogeneous*. The direct summands

$$A_{i,l} := p_{i,l} (C(X_{i,l}) \otimes \mathcal{K}) p_{i,l}$$

of A_i are called *homogeneous*. The spaces $X_{i,l}$ may always be assumed to be connected and have finite covering dimension by [11], and we make these assumptions from here on. We refer to the sequence (A_i, ϕ_i) as an *AH sequence*.

Now let A be a unital AH algebra. If A is the limit of an AH sequence (A_i, ϕ_i) for which

$$\liminf_{i \rightarrow \infty} \max_{1 \leq l \leq n_i} \frac{\dim(X_{i,l})}{\text{rank}(p_{i,l})} = 0,$$

then we say that A has *slow dimension growth*; if it is the limit of an AH sequence such that for some $M > 0$, we have $\dim(X_{i,l}) < M$, then we say that A has *no dimension growth*.

Given an AH sequence (A_i, ϕ_i) and $j > i$, we write $\phi_{i,j}$ for the composition $\phi_{j-1} \circ \cdots \circ \phi_i$ and $\phi_{i,\infty}$ for the canonical map from A_i into the limit algebra A . We define $\phi_{i,j}^{l,k} : A_{i,l} \rightarrow A_{j,k}$ and $\phi_{i,j}^k : A_i \rightarrow A_{j,k}$ to be the obvious restrictions of $\phi_{i,j}$. The $\phi_{i,j}^{l,k}$ are referred to as *partial maps*.

It is well known that an AH algebra $A = \lim_{i \rightarrow \infty} (A_i, \phi_i)$ is simple if and only if for every $i \in \mathbb{N}$ and $a \in A_i \setminus \{0\}$, there is some $j \geq i$ such that $\phi_{i,j}(a)$ generates A_j as an ideal. This last condition is equivalent to $\phi_{i,j}(a)$ being nonzero at every $x \in X_{j,1} \cup \cdots \cup X_{j,n_j}$.

2.3. Maps between homogeneous C^* -algebras. Let X and Y be compact connected metric spaces, and let $p \in C(X) \otimes \mathcal{K}$ and $q \in C(Y) \otimes \mathcal{K}$ be projections. Let

$$ev_x : p(C(X) \otimes \mathcal{K})p \rightarrow M_{\text{rank}(p)}$$

be given by $f \mapsto f(x)$; define ev_y for $y \in Y$ similarly. Let

$$\phi : p(C(X) \otimes \mathcal{K})p \rightarrow q(C(Y) \otimes \mathcal{K})q$$

be a unital $*$ -homomorphism. It is well known that for any $y \in Y$, the map $ev_y \circ \phi$ has the following form, up to unitary equivalence:

$$ev_y \circ \phi = \bigoplus_{j=1}^{N:=\frac{\text{rank}(q)}{\text{rank}(p)}} ev_{x_j},$$

where the x_j are points in X , not necessarily distinct. In other words, the x_j form an N – multiset, which we denote by $\text{sp}_\phi(y)$. The set of all such multisets is referred to as the N^{th} symmetric power of X , and is denoted by $P^N X$; it may be identified with the quotient of the Cartesian product X^N by the action of the symmetric group S_N on co-ordinates, and so inherits naturally a metric from X .

2.4. Semicontinuous projection-valued maps. Let X be a connected topological space. By a lower semicontinuous function $f : X \rightarrow M_n(\mathbb{C})_+$ we will mean a function such that for every vector $\xi \in \mathbb{C}^n$, the real-valued function $x \mapsto \langle f(x)\xi \mid \xi \rangle$ is lower semicontinuous (cf. [1]). The following result from [4], will be used in the sequel.

Proposition 2.1. *Let X be a compact metrisable Hausdorff space of dimension d , and let $P : X \rightarrow M_n(\mathbb{C})_+$ be a lower semicontinuous projection-valued map. Suppose that*

$$\text{rank}(P(x)) > \frac{1}{2}(d+1) + k, \forall x \in X.$$

It follows that there is a continuous projection-valued map $R : X \rightarrow M_n$ of constant rank equal to k such that

$$R(x) \leq P(x), \forall x \in X.$$

Remark 2.2. If we replace $(1/2)(d+1)$ with $d+1$ in the hypotheses of Proposition 2.1, then we may assume that the projection-valued map R corresponds to a trivial complex vector bundle over X . This is a consequence of the stability properties of vector bundles.

Lemma 2.3. *Let X, Y, p, q , and $\phi : p(C(X) \otimes \mathcal{K})p \rightarrow q(C(Y) \otimes \mathcal{K})q \subset M_m(C(Y))$ be as in Subsection 2.3 with X and Y not necessarily connected. Let O be an open subset of X , and let $r \in p(C(X) \otimes \mathcal{K})p$ be a positive element which is equal to a projection at every $x \in O$. Define a projection-valued map $R : Y \rightarrow M_m(\mathbb{C})$ as follows: $R(y)$ is the image of r under the direct sum of those irreducible direct summands of $ev_y \circ \phi$ which correspond to points in O . It follows that R is lower semicontinuous.*

Proof. For any $y \in Y$, let E_y denote the submultiset of $\text{sp}_\phi(y)$ consisting of those points which lie in O . Fix $y_0 \in Y$, and let δ denote the smallest distance between a point in E_{y_0} and a point in the complement of O . The map $y \mapsto \text{sp}_\phi(y)$ is continuous, whence there is an open

neighbourhood V of y_0 such that, for each $y \in V$, the submultiset F_y of $\text{sp}_\phi(y)$ consisting of those points which are at distance at most $\delta/2$ from some point in E_{y_0} has the same cardinality as E_{y_0} , and moreover the map $y \mapsto F_y$ is continuous.

Define a continuous projection-valued map $\tilde{R} : V \rightarrow M_m(\mathbb{C})$ as follows: $\tilde{R}(y)$ is the image of r under the sum of the irreducible direct summands of $ev_y \circ \phi$ which correspond to the elements of F_y . We have that $\tilde{R}(y) \leq R(y)$ for every $y \in V$, and that $\tilde{R}(y_0) = R(y_0)$. Let $z_n \rightarrow y_0$. For all n sufficiently large we have $z_n \in V$, whence, for each $\xi \in \mathbb{C}^{\text{rank}(q)}$, we have $\langle R(z_n)\xi, \xi \rangle \geq \langle \tilde{R}(z_n)\xi, \xi \rangle$. It follows that

$$\liminf_{n \rightarrow \infty} \langle R(z_n)\xi, \xi \rangle \geq \lim_{n \rightarrow \infty} \langle \tilde{R}(z_n)\xi, \xi \rangle = \langle \tilde{R}(y_0)\xi, \xi \rangle = \langle R(y_0)\xi, \xi \rangle,$$

and so R is lower semicontinuous. The assumption that Y is connected is not really needed. $\square$

3. A WORD ON STRATEGY

Before plunging headlong into the technical details of our proof, let us attempt to explain why a unital simple AH algebra with no dimension growth *ought* to absorb the Jiang-Su algebra tensorially.

Let $p, q \geq 2$ be relatively prime integers. Bearing in mind the isomorphism $M_{pq} \cong M_p \otimes M_q$, one defines

$$I_{p,q} = \{f \in C([0, 1]; M_{pq}) \mid f(0) \in \mathbf{1}_p \otimes M_q, f(1) \in M_p \otimes \mathbf{1}_q\}.$$

The algebra $I_{p,q}$ is referred to as a *prime dimension drop algebra*, and the Jiang-Su algebra, denoted by $\mathcal{Z}$, is the unique unital simple inductive limit of prime dimension drop algebras with the same K-theory and tracial state space as the algebra of complex numbers (see [12]). In order to prove that a unital C^* -algebra absorbs the Jiang-Su algebra tensorially, it suffices to prove that for each p, q as above, there is an approximately central sequence of unital $*$ -homomorphisms $\gamma_n : I_{p,q} \rightarrow A$ (cf. [14, Proposition 2.2]).

Let $A = \lim_{i \rightarrow \infty} (A_i, \phi_i)$ be a unital simple AH algebra with no dimension growth, and assume for simplicity that each A_i is homogeneous with connected spectrum X_i . Fix a finite subset F of A_i . It is known that for any $\epsilon > 0$ there exists $j > i$ such that for every $y \in X_j$, the finite-dimensional representation $ev_y \circ \phi_{i,j}$ of A_i has the following property: the multiset $\text{sp}_{\phi_{i,j}}(y)$ can be partitioned into submultisets $S_1, \dots, S_m$ such that (a) all of the elements in a fixed S_t lie in a ball of radius at most ϵ , and (b) each S_t has large cardinality relative to $\dim(X_j)$. Suppose that $S_t = \{\{x_1, \dots, x_k\}\}$ (the double brackets indicate a multiset, i.e., a set with repetition). The projections $ev_{x_1}(\mathbf{1}_{A_i}), \dots, ev_{x_k}(\mathbf{1}_{A_i})$ (whose sum is denoted by I_t) are pairwise orthogonal and Murray-von Neumann equivalent, and so they and the partial isometries implementing the said equivalences generate a copy of M_k which almost commutes with the image of F under the map $I_t(ev_y \circ \phi_{i,j})I_t$. If k is large enough, then there is a unital $*$ -homomorphism from $I_{p,q}$ into M_k which almost commutes with the image of F . Repeating this procedure for each of $S_1, \dots, S_m$, we obtain a unital $*$ -homomorphism from $I_{p,q}$ into the fibre $M_{\text{rank}(\mathbf{1}_{A_j})}$ of A_j over $y \in X_j$ which almost commutes with the image of F . By the semiprojectivity of $I_{p,q}$, this $*$ -homomorphism can be extended to have codomain equal to the restriction of A_j to a closed neighbourhood of y . Thus, it is straightforward to

see the existence of the required γ_n s in a “local” sense. This article handles the passage from local to global. What makes this possible is the fact that the homotopy groups of the space of k -dimensional representations of $I_{p,q}$ vanish in low dimensions (cf. [5]).

4. EXCISING POINT EVALUATIONS

Let A be a unital simple AH algebra with slow dimension growth. We say that an AH sequence (A_i, ϕ_i) with limit A *realises slow dimension growth* if

$$\liminf_{i \rightarrow \infty} \max_{1 \leq l \leq n_i} \frac{\dim(X_{i,l})}{\text{rank}(p_{i,l})} = 0;$$

assume that (A_i, ϕ_i) is such a sequence. Our goal in this section is to prove that for each finite subset F of A_i , there is some $j > i$ with the property that the bonding map $\phi_{i,j}$ is “almost” a direct sum of a suitably dense family of irreducible representations of A_i together with a second map $\overline{\phi_{i,j}}$.

Let X be a compact metric space and $N \geq 1$ an integer. Let $O_1, \dots, O_m$ be open subsets of X whose closures are pairwise disjoint. The C^* -subalgebra of $A = M_N(C(X))$ consisting of those functions $f : X \rightarrow M_N(\mathbb{C})$ that are constant on each O_s is denoted by $A_{\{O_1, \dots, O_m\}}$. It is easily verified that $A_{\{O_1, \dots, O_m\}} \cong M_N(C(X'))$ where X' is the quotient of X obtained by shrinking each set $\overline{O_s}$ to a distinct point w_s , $s = 1, \dots, m$.

If $\rho : A \rightarrow B$ is a $*$ -homomorphism we will write $L \cdot \rho$ for a $*$ -homomorphism $A \rightarrow M_L(B)$ which is unitarily equivalent to the direct sum of L copies of ρ .

Lemma 4.1. *Let X, Y be compact metrisable spaces and let*

$$\gamma : A = M_N(C(X)) \rightarrow q(C(Y) \otimes \mathcal{K})q$$

be a unital $$ -homomorphism. Let $\{O_s\}_{s=1}^m$ be open subsets of X with disjoint closures. Suppose that for all $s = 1, \dots, m$ and all $y \in Y$*

$$|\text{sp}_\gamma(y) \cap O_s| \geq (K+2)\dim(Y).$$

Then the restriction of γ to $A_{\{O_1, \dots, O_m\}}$ decomposes as a direct sum $\bar{\gamma} \oplus L \cdot \rho$ where $L \geq K\dim(Y)$ and ρ is $$ -homomorphism with finite dimensional image and spectrum equal to $\{w_1, \dots, w_m\}$.*

Proof. We may assume that $q \in M_R(C(Y))$ for some $R \geq 1$. Fix a system of matrix units (p_{cd}) for $M_N(\mathbb{C})$. For each $y \in Y$, let $q_{cd}^{(s)}(y)$ be the image of p_{cd} under the direct sum of all the irreducible direct summands of $ev_y \circ \gamma$ which correspond to points in O_s . By Lemma 2.3 we see that $q_{11}^{(s)}(y)$ is a lower semicontinuous projection-valued map on Y whose rank is at least $(K+2)\dim(Y)$ at every point.

Apply Proposition 2.1 and the remark thereafter to find a continuous constant rank subprojection $r_{11}^{(s)} : Y \rightarrow M_R(\mathbb{C})$ of $q_{11}^{(s)}$ whose rank L is at least $K\dim(Y)$ and which corresponds to a trivial vector bundle over Y . Since $r_{11}^{(s)}(y) \leq q(y)$ for all $y \in Y$ it follows that $r_{11}^{(s)} \in q(C(Y) \otimes \mathcal{K})q$. Set $r_{cd}^{(s)} = q_{c1}^{(s)} r_{11}^{(s)} q_{1d}^{(s)} = \gamma(p_{c1}) r_{11}^{(s)} \gamma(p_{1d})$. It is straightforward to check that $\{r_{cd}^{(s)}\}_{c,d=1}^N$ is a system of matrix units in $q(C(Y) \otimes \mathcal{K})q$. Let I_s denote the unit of the subalgebra of $q(C(Y) \otimes \mathcal{K})q$ generated by the $r_{cd}^{(s)}$.

To complete the proof of the Lemma, it will suffice to show that, up to unitary equivalence,

$$I_s(\gamma|_{A_{\{O_1, \dots, O_m\}}})I_s = \bigoplus_{t=1}^L ev_{w_s} = L \cdot ev_{w_s}.$$

We must first show that the left hand side is a $*$ -homomorphism. Fix $y \in Y$. Observe that the irreducible direct summands of $ev_y \circ \gamma$ which correspond to points in O_s are, upon restricting γ to $A_{\{O_1, \dots, O_s\}}$, replaced by irreducible representations of $A_{\{O_1, \dots, O_s\}}$ corresponding to the point $w_s \in \text{Spec}(A_{\{O_1, \dots, O_s\}})$. In particular, the image of any $a \in A_{\{O_1, \dots, O_s\}}$ under these irreducible representations is contained in the linear span of the $q_{cd}^{(s)}(y)$, and so commutes with I_s (an easy exercise using the definition of the $r_{cd}^{(s)}$ shows that $I_s q_{cd}^{(s)} = q_{cd}^{(s)} I_s = r_{cd}^{(s)}$). Since I_s commutes with the image of $\gamma|_{A_{\{O_1, \dots, O_m\}}}$, we see that $I_s(\gamma|_{A_{\{O_1, \dots, O_m\}}})I_s$ is a $*$ -homomorphism.

The map $I_s(\gamma|_{A_{\{O_1, \dots, O_m\}}})I_s$ factors through the evaluation of $A_{\{O_1, \dots, O_s\}}$ at w_s , and has multiplicity L . To see that this finite-dimensional representation of $A_{\{O_1, \dots, O_s\}}$ decomposes as the direct sum of L representations of multiplicity one, we observe that $r_{11}^{(s)}$ can be decomposed into the direct sum of L equivalent rank one projections by virtue of its triviality. Let ξ be one such projection. We can form matrix units $\xi_{cd} = q_{c1}^{(s)} \xi q_{1d}^{(s)}$ to obtain an irreducible subrepresentation of $I_s(\gamma|_{A_{\{O_1, \dots, O_m\}}})I_s$ of multiplicity one. There are L such subrepresentations, and they are mutually orthogonal. This completes the proof of the Lemma. $\square$

Lemma 4.2. *Let A be an infinite dimensional unital simple AH algebra with slow dimension growth, and let (A_j, ϕ_j) be an AH sequence which realises the slow dimension growth of A . Suppose that $A_i = M_N(C(X_i))$ for some i and let there be given $F \subseteq A_i$ finite, a tolerance $\epsilon > 0$, a natural number K , and a finite set $\{x_1, \dots, x_m\} \subseteq X_i$.*

It follows that there are $j > i$, open neighborhoods O_s of x_s ($s = 1, \dots, m$) in X_i with pairwise disjoint closures, and a finite set $F' \subseteq A'_i := A_{i\{O_1, \dots, O_m\}}$ with the following properties for each $k \in \{1, \dots, n_j\}$:

- (i) $F' \approx_\epsilon F$;
- (ii) *The map $\gamma_{j,k} : A'_i \rightarrow A_{j,k}$ obtained by restricting $\phi_{i,j}^k$ to A'_i is, up to unitary equivalence inside its codomain, of the form $\overline{\gamma} \oplus L \cdot \rho$, where $L \in \mathbb{N}$, ρ is a $*$ -homomorphism with finite dimensional image and spectrum consisting of the points $\{w_1, \dots, w_m\}$ corresponding to the images of the sets $\overline{O_1}, \dots, \overline{O_m}$ in the quotient space of X_i representing the spectrum of A'_i ;*
- (iii) $L \geq K \dim(X_{j,k})$.

Proof. Part (i) of the conclusion of the lemma follows from a standard approximation argument. We will show that the associated choice of $O_1, \dots, O_m$ suffices for the conclusion of the lemma proper. Set $\gamma_{j,k} = \phi_{i,j}^k$ with i fixed. For $j \geq i$, let $L(j, k, s)(y)$ denote the number of irreducible direct summands which correspond to points in O_s of the finite-dimensional representation $ev_y \circ \gamma_{j,k}$ of A_i , where $y \in X_{j,k}$, $k \in \{1, \dots, n_j\}$ and $s \in \{1, \dots, m\}$. Thus $L(j, k, s)(y) = |\text{sp}_{\gamma_{j,k}}(y) \cap O_s|$.

Choose positive elements $a_1, \dots, a_m \in A_i$ such that $\text{supp}(a_s) = O_s$. By the simplicity of A , there exist $j_0 > i$ and $M \geq 1$ such that for each $j \geq j_0$, $s \in \{1, \dots, m\}$ and $k \in \{1, \dots, n_j\}$,

there are elements $b_1, \dots, b_M$ in $A_{j,k}$ such that $\sum_{t=1}^M b_t \gamma_{j,k}(a_s) b_t^* = p_{j,k}$. It follows that for each $y \in X_{j,k}$

$$M \cdot N \cdot L(j, k, s)(y) = M \cdot \text{rank}(\gamma_{j,k}(a_s)(y)) \geq \text{rank}(p_{j,k}(y))$$

and hence

$$\frac{\dim(X_{j,k})}{L(j, k, s)(y)} \leq \frac{\dim(X_{j,k})}{\text{rank}(p_{j,k})} \cdot M \cdot N.$$

By the slow growth dimension condition if j is large enough then $L(j, k, s)(y) \geq (K + 2)\dim(X_{j,k})$ for all s, k and all $y \in X_{j,k}$. Properties (ii) and (iii) in the conclusion of the present lemma now follow from an application of Lemma 4.1. $\square$

5. APPROXIMATE RELATIVE COMMUTANTS

The homogeneous C^* -algebras considered in this section are not necessarily of the form $qM_N(C(Y))q$, but only locally isomorphic to such algebras. The goal of this section is to prove Proposition 5.4.

Let X be a compact metrisable space. Let $\epsilon > 0$ and a finite set $F \subseteq C(X)$ be given. Let R be a finite subset of X . Let $\gamma : C(X) \rightarrow B$ be a unital $*$ -homomorphism, where B is a unital separable homogeneous C^* -algebra with spectrum Y . Given an integer $K \geq 1$, we say that γ admits a K -large system of compatible local finite dimensional approximations with respect to the data ϵ, F and R if there are two finite closed covers $\{W_s\}_{s=1}^M$ and $\{V_s\}_{s=1}^M$ of Y with $W_s \subset \overset{\circ}{V}_s$ such that for each s there is a partition of unity of B into projections $e(s, i)$ defined on V_s ,

$$\sum_{i=1}^{n(s)} e(s, i) = 1_B|_{V_s}$$

with the following properties.

(i) The restrictions of $e(s_1, i_1), \dots, e(s_m, i_m)$ to any nonempty intersection $V_{s_1} \cap \dots \cap V_{s_m} \neq \emptyset$ mutually commute and the rank of the product $e(s_1, i_1) \cdot \dots \cdot e(s_m, i_m)$ is either 0 or $\geq K$ at all points of $V_{s_1} \cap \dots \cap V_{s_m}$.

(ii) For any s there are points $x_{\kappa(s,1)}, \dots, x_{\kappa(s,n(s))}$ in R such that for all $f \in F$

$$\|\gamma(f) - \sum_{i=1}^{n(s)} f(x_{\kappa(s,i)}) e(s, i)\|_{V_s} < \epsilon/2.$$

Let X be a compact metrisable space. For $\delta > 0$ we denote by $r(X, \delta)$ the smallest number r with the property that for every finite set $G \subset X$ there are open subsets $O_1, \dots, O_r$ of X of diameter $< \delta$ whose union contains G and such that their closures are mutually disjoint. One can see that $r(X, \delta) < \infty$ by embedding X in the Hilbert cube and choosing the O_s to be parallelepipeds of the form

$$X \cap \left(\prod_{i \leq N} (a_i, b_i) \times \prod_{i > N} [0, 1] \right).$$

If $F \subset pM_N(C(X))p \subset M_N(C(X))$ is a finite subset we denote $\omega(F, \delta)$ the δ -oscillation of the family F :

$$\omega(F, \delta) = \sup\{\|f(x) - f(x')\|_{M_N(\mathbb{C})} : d(x, x') < \delta\}$$

Proposition 5.1. *Let X be a compact metric space and let $\gamma : C(X) \rightarrow B$ be a unital $*$ -homomorphism to a separable homogeneous C^* -algebra B with spectrum of dimension d . Let $\delta > 0$ and suppose that γ admits a direct sum decomposition of the form $\gamma = \phi \oplus L \cdot \rho$ where ρ is a $*$ -homomorphism with finite dimensional image whose spectrum R is δ -dense in X and $L \geq ((r(X, \delta) + 1)^{d+1} - 1)K$. If $F \subset C(X)$ is a finite set then γ admits a K -large system of compatible local finite dimensional approximations with respect to the data $\epsilon = 2\omega(F, 3\delta)$, F and R .*

Proof. Suppose that γ, ϕ, ρ and L are as in the statement. The cover $\{V_s\}_s$ and the corresponding partitions of unity are constructed as follows. Set $r = r(X, \delta)$. By the compactness of Y there is a finite open cover $\mathcal{V} = \{V_1, \dots, V_M\}$ of the spectrum Y of B such that for each V_s there is a family $O_{(s,1)}, \dots, O_{(s,r)}$ of open subsets of X of diameter $< \delta$, whose union contains $\text{sp}_\phi(y)$ for all $y \in V_s$ and such that $\overline{O_{(s,i)}} \cap \overline{O_{(s,j)}} = \emptyset$ for $i \neq j$. Since $\dim(Y) = d$, after passing to a finer subcover of $\mathcal{V}$, we may arrange that there is an open cover $\{V_1, \dots, V_M\}$ of Y which can be colored in $(d+1)$ -colors such that the elements of the same color have disjoint closures. In other words we can write $\{V_1, \dots, V_M\}$ as a disjoint union $\mathcal{V}_1 \cup \dots \cup \mathcal{V}_{d+1}$ such that if $V_s, V_t \in \mathcal{V}_i$ for some $1 \leq i \leq d+1$, and $s \neq t$, then $\overline{V_s} \cap \overline{V_t} = \emptyset$. Let us note that by enlarging each V_s to a set of the form $\{y \in Y : d(y, V_s) \leq \alpha\}$ we may arrange that in addition to the above properties, each V_s is closed and its interior contains some closed subset W_s such that $\{W_s\}_{s=1}^M$ is a cover of Y .

We need to work with the coloring map $\{1, \dots, M\} \rightarrow \{1, \dots, d+1\}$, $s \mapsto \bar{s}$ where $\bar{s}$ is defined by the condition that V_s has color $\bar{s}$, i.e., V_s is an element of the family $\mathcal{V}_{\bar{s}}$. Consider the set S consisting of all sets $a = \{(s_1, i_1), \dots, (s_m, i_m)\}$ with the property that

$$V_{s_1} \cap \dots \cap V_{s_m} \neq \emptyset, \quad O_a := O_{(s_1, i_1)} \cap \dots \cap O_{(s_m, i_m)} \neq \emptyset,$$

where $1 \leq s_1, \dots, s_m \leq M$ are mutually distinct (hence necessarily $m \leq d+1$ since distinct sets V_s of the same color are disjoint) and $1 \leq i_1, \dots, i_m \leq r$. Consider also the set $\hat{S}$ consisting of all sets of the form $\{(\bar{s}_1, i_1), \dots, (\bar{s}_m, i_m)\}$ where $m \leq d+1$, $\bar{s}_1, \dots, \bar{s}_m$ are mutually distinct elements (colors) in the set $\{1, \dots, d+1\}$ and $1 \leq i_1, \dots, i_m \leq r$. Note that $\hat{S}$ has $(1+r)^{d+1} - 1$ elements and observe that $\hat{S} \supset \{\bar{a} : a \in S\}$ where for $a \in S$ as above, we set $\bar{a} = \{(\bar{s}_1, i_1), \dots, (\bar{s}_m, i_m)\}$.

By replacing ϕ by $\phi \oplus (L - L_0) \cdot \rho$, we may assume that $L = L_0$ where $L_0 = ((1+r)^{d+1} - 1)K$. Then we can identify $L \cdot \rho$ with the $*$ -homomorphism

$$\sigma : C(X) \rightarrow qBq \otimes \mathcal{L}(\ell^2(\hat{S})) \otimes M_K(\mathbb{C})$$

$$\sigma(f) = \rho(f) \otimes 1 \otimes 1_K.$$

By assumption, $\rho : C(X) \rightarrow qBq$, $q = \rho(1)$, must have the form

$$\rho(f) = \sum_{k=1}^c f(x_k) q_k$$

where $q_1 + \dots + q_c = q$ are mutually orthogonal nonzero projections and the spectrum $R = \{x_1, \dots, x_c\}$ of ρ is such that for any $x \in X$ there is $x_k \in R$ such that $d(x, x_k) < \delta$. Therefore there is a map $\kappa : S \rightarrow \{1, \dots, c\}$ with the property that

$$(1) \quad d(x_{\kappa(a)}, O_a) < \delta.$$

For each fixed open set V_s we are going to define a partition of $1_B|_{V_s}$. For $1 \leq i \leq r$, let $h_{(s,i)}$ be an element of $C(X)$ such that $h_{(s,i)}(x) = 1$ for all $x \in O_{(s,i)}$ and such that $h_{(s,i)}(x) = 0$ on $\bigcup_{j \neq i} O_{(s,j)}$. Define open projections $p(s, i) = \phi(h_{(s,i)})|_{V_s}$ for $1 \leq i \leq r$. For each (s, i) let $S(s, i)$ be the subset of those elements $a \in S$ with the property that $(s, i) \in a$. Let $T = \{1, \dots, c\} \times \hat{S}$ and set

$$T(s, i) = \{(\kappa(a), \bar{a}) : a \in S(s, i)\} \subset T.$$

Let $\xi(s, r+1), \dots, \xi(s, n(s))$ be an enumeration of the complement set of $\bigcup_{i=1}^r T(s, i)$ in T . If this complement set is nonempty then $n(s) > r$; otherwise set $n(s) = r$. Set $T(s, i) = \{\xi(s, i)\}$ for $i = r+1, \dots, n(s)$ and let us observe that for each s the family $(T(s, i))_{i=1}^{n(s)}$ forms a partition of T . Indeed, for $1 \leq i \neq j \leq r$, $T(s, i) \cap T(s, j) = \emptyset$ since for each $a = \{(s_1, i_1), \dots, (s_m, i_m)\} \in S$, the colours $\bar{s}_1, \dots, \bar{s}_r$ are mutually distinct, because $V_{s_1} \cap \dots \cap V_{s_m} \neq \emptyset$.

After this preparation, for each $(s, i) \in T$ we define a projection

$$q(s, i) = \sum_{(k,b) \in T(s,i)} q_k \otimes \chi_{\{b\}} \otimes 1_K \in qBq \otimes \mathcal{L}(\ell^2(\hat{S})) \otimes M_K(\mathbb{C})$$

(recall that $q_1, \dots, q_c$ are the spectral projection of ρ). We also define projections $(e(s, i))_{i=1}^{n(s)}$ on V_s by

$$e(s, i) = \begin{cases} p(s, i) + q(s, i), & \text{if } 1 \leq i \leq r \\ q(s, i), & \text{if } r < i \leq n(s). \end{cases}$$

Then

$$\sum_{i=1}^{n(s)} e(s, i) = 1_B|_{V_s}$$

is a partition of unity on V_s . Indeed $\sum_{i=1}^r p(s, i) = \sum_{i=1}^r \phi(h_{(s,i)}) = \phi(1)|_{V_s}$ and $\sum_{i=1}^{n(s)} q(s, i) = \sigma(1)|_{V_s}$ since $(T(s, i))_{i=1}^{n(s)}$ is a partition of T . Note that if $O(s, i) = \emptyset$ then $S(s, i) = T(s, i) = \emptyset$ and $p(s, i) = q(s, i) = 0$.

It remains to verify the properties (i) and (ii). To verify condition (i) we observe first that for each $a = \{(s_1, i_1), \dots, (s_m, i_m)\} \in S$ the rank of the product $e(s_1, i_1) \cdot \dots \cdot e(s_m, i_m)$ is either 0 or $\geq K$. Indeed if all indices $i_1, \dots, i_m$ are $\leq r$, then

$$e(s_1, i_1) \cdot \dots \cdot e(s_m, i_m) \geq q(s_1, i_1) \cdot \dots \cdot q(s_m, i_m) \geq q_{\kappa(a)} \otimes \chi_{\{\bar{a}\}} \otimes 1_K$$

which follows simply because $a \in S(s_1, i_1) \cap \dots \cap S(s_m, i_m)$. If $i_k > r$ for some k , then the rank of $e(s_k, i_k)$ is divisible by K and hence so is the rank of the product.

Second, if $V_{s_1} \cap \dots \cap V_{s_m} \neq \emptyset$ then the projections $e(s_1, i_1), \dots, e(s_m, i_m)$ commute on $V_{s_1} \cap \dots \cap V_{s_m}$ by construction.

Let us now verify property (ii) for a fixed V_s . The number $\kappa(\{(s, i)\})$, which we will write from now on as $\kappa(s, i)$, was defined whenever $\{(s, i)\} \in S$. It is convenient to extend this

notation as follows. If $1 \leq i \leq r$ but $\{(s, i)\} \notin S$ set $\kappa(s, i) = 1$ and if $i > r$ we let $\kappa(s, i)$ denote the (first) coordinate of $\xi(s, i)$ in $\{1, \dots, c\}$. We are going to show that if $f \in F$, then

$$(2) \quad \|\gamma(f) - \sum_{i=1}^{n(s)} f(x_{\kappa(s,i)})e(s, i)\|_{V_s} \leq \epsilon/2.$$

Define $\phi'_s(f)(y) = \sum_{i=1}^r f(x_{\kappa(s,i)})p(s, i)(y)$ for $y \in V_s$ and

$$\sigma'_s(f) = \sum_{i=1}^r f(x_{\kappa(s,i)})q(s, i) + \sum_{i=r+1}^{n(s)} f(x_{\kappa(s,i)})e(s, i).$$

Recall that for $y \in V_s$, $\phi(f)(y)$ depends only on the restriction of f to $\bigcup_{i=1}^r O_{(s,i)}$. Since $d(x_{\kappa(s,i)}, O_{(s,i)}) < \delta$ for $1 \leq i \leq r$ and $|f(x) - f(x')| \leq \epsilon/2$ if $d(x, x') < 3\delta$ and $f \in F$ it follows that

$$(3) \quad \|\phi(f) - \phi'_s(f)\|_{V_s} \leq \epsilon/2, \quad \forall f \in F.$$

Since T is partitioned into the sets $(T(s, i))_{i=1}^r$ and $\{\xi(s, i)\}, i = r+1, \dots, n(s)$ we can write

$$(4) \quad \sigma(f) = \sum_{i=1}^r \sum_{(k,b) \in T(s,i)} f(x_k) q_k \otimes \chi_{\{b\}} \otimes 1_K + \sum_{i=r+1}^{n(s)} f(x_{\kappa(s,i)})e(s, i).$$

Note that if $(k, b) \in T(s, i)$ for $1 \leq i \leq r$ then $(k, b) = (\kappa(a), \bar{a})$ for some $a \in S(s, i)$. On the other hand if $a \in S(s, i)$, we see that $d(x_{\kappa(a)}, x_{\kappa(s,i)}) < 3\delta$ using (1) and the inclusion $O_a \subset O_{(s,i)}$. Since $q(s, i) = \sum_{(k,b) \in T(s,i)} q_k \otimes \chi_{\{b\}} \otimes 1_K$, eq. (4) leads to

$$(5) \quad \|\sigma(f) - \sigma'_s(f)\| \leq \epsilon/2, \quad \forall f \in F.$$

Let us set $\gamma'_s = \phi'_s \oplus \sigma'_s$. Recalling that $\gamma = \phi \oplus \sigma$ we then obtain

$$(6) \quad \|\gamma(f) - \gamma'_s(f)\|_{V_s} \leq \epsilon/2, \quad \forall f \in F.$$

This completes the proof of (2) since $\gamma'_s(f) = \sum_{i=1}^{n(s)} f(x_{\kappa(s,i)})e(s, i)$. $\square$

Under the same assumptions as in Proposition 5.1 we establish two Lemmas. Consider the C^* -algebra $B^\sharp$ consisting of those elements g of B which commute with all projections $e(s, i)$, $i = 1, \dots, n(s)$ on each closed set W_s .

Lemma 5.2. *If $g \in B^\sharp$ and $\|g\| \leq 1$, then $\|[g, \gamma(f)]\| \leq \epsilon$ for all $f \in F$.*

Proof. If $f \in F$, then g commutes with $\gamma'_s(f)$ on $W_s \subset V_s$ and hence

$$\|[\gamma(f), g]\| \leq \sup_s \|[\gamma(f) - \gamma'_s(f), g]\|_{W_s} \leq 2 \sup_s \|g\| \|\gamma(f) - \gamma'_s(f)\|_{W_s} \leq \epsilon,$$

by using (6). $\square$

Lemma 5.3. *$B^\sharp$ is a unital separable continuous field C^* -algebra over Y whose fibers have all their irreducible representations of dimension $\geq K$.*

Proof. Let us note that $B^\sharp$ is a $C(Y)$ -subalgebra of a continuous field B of matrices and hence $B^\sharp$ is a continuous field C^* -algebra itself. For $y \in Y$, let $\pi_y : B \rightarrow B(y)$ be the evaluation map. For each fixed y find a maximal set of indices $\{s_1, \dots, s_m\}$ such that $y \in W_{s_1} \cap \dots \cap W_{s_m}$. Then y has a neighborhood $V \subset V_{s_1} \cap \dots \cap V_{s_m}$ such that $V \cap W_s = \emptyset$ for all $s \notin \{s_1, \dots, s_m\}$. Let S_y consist of all elements a of S of the form $a = \{(s_1, i_1), \dots, (s_m, i_m)\}$. For each $a \in S_y$ let $b(a)$ be an arbitrary element of B which vanishes outside V and set $e(a) = e(s_1, i_1) \cdot \dots \cdot e(s_m, i_m)$. Then

$$b^\sharp = \sum_{a \in S_y} e(a)b(a)e(a)$$

commutes with all $e(s_k, i_k)$ on W_{s_k} , $1 \leq k \leq m$ and vanishes on all W_s with $s \notin \{s_1, \dots, s_m\}$. This shows that $b^\sharp \in B^\sharp$. Since $e(a) \cdot e(b) = 0$ if $a \neq b$ and since $\pi_y(b(a))$ can be chosen to be any element of $B(y)$ we see that

$$\pi_y(B^\sharp) \cong \bigoplus_a M_{r(a)}(\mathbb{C})$$

where a runs in S_y and $r(a) = \text{rank}(e(a))$ is either 0 or $\geq K$. $\square$

Proposition 5.4. *Given I_{p_1, p_2} there is an integer $\ell \geq 1$ with the following property. Let X be a compact metric space and let $\gamma : M_N(C(X)) \rightarrow B$ be a unital $*$ -homomorphism to a separable homogeneous C^* -algebra B with spectrum of dimension d . Let $\delta > 0$ and suppose that γ decomposes as a direct sum $\gamma = \phi \oplus L^{d+2} \cdot \rho$ where ρ is a $*$ -homomorphism with finite dimensional image whose spectrum is δ -dense in X and such that $L \geq r(X, \delta) + 1 + \ell$. Then there is a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow B$ such that $\|[\eta(g), \gamma(f)]\| \leq 2\omega(F, 3\delta)$ for all $g \in I_{p_1, p_2}$, $\|g\| \leq 1$ and $f \in F$.*

Proof. In the first part of the proof we consider the case $N = 1$. By [5, Theorem 6.2] there is ℓ depending only on p_1, p_2 with the following property. If D is a separable recursive subhomogeneous algebra of finite topological dimension d and minimum matrix size $\geq \ell(d + 1)$, then there is a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow D$.

Let $\delta > 0$ be given and set $r = r(X, \delta)$ and suppose that γ is as in the statement with $L \geq r + 1 + \ell$. Then $L^{d+2} \geq (d + 2)(r + 1)^{d+1}\ell \geq ((r + 1)^{d+1} - 1)K$ where $K = \ell(d + 1)$. By Proposition 5.1, γ admits a K -large system of compatible local finite dimensional approximations and moreover by Lemma 5.3 the corresponding commutant C^* -algebra $B^\sharp$ is a unital separable continuous field with fibers finite dimensional C^* -algebras whose all direct summands have size $\geq K$. It follows from [2, Theorem 4.6] that there is a recursive subhomogeneous algebra D of finite topological dimension d and minimum matrix size $\geq \ell(d + 1)$ such that $D \subset B^\sharp$. By [5, Theorem 6.2] there is a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow D$. By Lemma 5.2 we conclude that $\|[\eta(g), \gamma(f)]\| \leq 2\omega(F, 3\delta)$ for all $g \in I_{p_1, p_2}$, $\|g\| \leq 1$ and $f \in F$.

Consider now the general case with $\gamma : M_N(C(X)) \rightarrow B$. Without any loss of generality we may assume that F is the union of a system of matrix units $(e_{\alpha, \beta})$ for $M_N(\mathbb{C}1_{C(X)})$ and a finite subset F_0 of $1_N \otimes C(X)$. Let ℓ be as above and $\delta > 0$. Let D be the commutant of $\gamma(M_N(\mathbb{C}))$ in B . Then D is a homogeneous C^* -algebra with spectrum Y and $\gamma \cong id_N \otimes \gamma_0 : M_N(\mathbb{C}) \otimes C(X) \rightarrow M_N(\mathbb{C}) \otimes D$ for some unital $*$ -homomorphism $\gamma_0 : C(X) \rightarrow D$. Moreover γ_0 can be written as a direct sum $\gamma_0 = \phi_0 \oplus L^{d+2} \cdot \rho_0$ where ρ_0 is a $*$ -homomorphism with

finite dimensional image whose spectrum is δ -dense in X . By the first part of the proof there is a unital $*$ -homomorphism $\eta_0 : I_{p_1, p_2} \rightarrow D$ such that $\|[\eta_0(g), \gamma_0(f)]\| \leq 2\omega(F_0, 3\delta)$ for all $g \in I_{p_1, p_2}$ and $f \in F_0$. Clearly $\eta := 1_N \otimes \eta_0$ has the desired properties since $\omega(F_0, 3\delta) = \omega(F, 3\delta)$. $\square$

Let (X, d) be a compact metric space and let $V_1, \dots, V_m$ be closed subsets of X . Let (X', d') be the compact metric space obtained by shrinking each V_s to a point w_s . Let $\pi : X \rightarrow X'$ be the quotient map. The induced metric d' on X' is given by

$$d'(\pi(x), \pi(y)) = \inf\{d(x_1, y_1) + \dots + d(x_n, y_n)\}$$

where the infimum is taken over all finite sequences $x_1, \dots, x_n$ and $y_1, \dots, y_n$ with $\pi(x) = \pi(x_1)$, $\pi(y) = \pi(y_n)$ and $\pi(y_i) = \pi(x_{i+1})$ for $i = 1, \dots, n-1$.

Let $A = \lim_{i \rightarrow \infty} (A_i, \phi_i)$, where each A_i is semi-homogeneous and assume that there is an $d \geq 0$ such that $\dim(\text{Spec}(A_i)) \leq d$ for every $i \in \mathbb{N}$.

Lemma 5.5. *Let A be as above and assume that $A_1 = M_N(C(X))$. For any finite subset $F \subset A_1$, any $\epsilon > 0$ and any relatively prime integers $p_1, p_2 \geq 2$ there are $j > 1$ and a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow A_j$ such that $\|[\eta(g), \phi_{1,j}(f)]\| \leq \epsilon$ for all $f \in F$ and $g \in I_{p_1, p_2}$, $\|g\| \leq 1$.*

Proof. Given F and ϵ , choose and fix $\delta > 0$ small enough so that $\omega_X(F, 4\delta) < \epsilon/10$. Let $\{x_1, \dots, x_m\}$ be a δ -dense subset of X and let $O_1, \dots, O_m$ be open sets in X with disjoint closures and such that $x_s \in O_s$ for $s = 1, \dots, m$. We may assume that these sets are sufficiently small so that there is a finite subset F' of $M_N(C(X))$ such that each $f' \in F'$ is constant on each O_s and $F' \approx_{\epsilon/10} F$. Moreover, by replacing the sets O_s by even smaller sets we may arrange that if (X', d') denotes the metric space obtained by shrinking each $\overline{O_s}$ to a point w_s , then $d(x, y) < 4\delta$ whenever $x, y \in X$ satisfy $d'(\pi(x), \pi(y)) < 3\delta$ where $\pi : X \rightarrow X'$ denotes the quotient map. Therefore $\omega_{X'}(F', 3\delta) \leq \omega_X(F', 4\delta) \leq \omega_X(F, 4\delta) + 2\epsilon/10 < \epsilon/3$. Let ℓ be given by Proposition 5.4. Applying Lemma 4.2 with $K = L^{d+2}$ (where $L \geq r(X', \delta) + 1 + \ell$) we find $j > 1$ such that the map $\gamma : M_N(C(X')) \rightarrow A_j$ obtained by restricting $\phi_{1,j}$ to $M_N(C(X'))$ decomposes as $\bar{\gamma} \oplus L^{d+2} \cdot \rho$ where ρ is a $*$ -homomorphism with finite dimensional image and whose spectrum is the set $\{w_1, \dots, w_m\}$ which is δ -dense in X' . Applying Proposition 5.4 we obtain a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow A_j \subset A$ such that $\|[\eta(g), \phi_{1,j}(f')]\| \leq 2\omega_{X'}(F', 3\delta) < 2\epsilon/3$ for all $f' \in F'$ and $g \in I_{p_1, p_2}$, $\|g\| \leq 1$. Since $F' \approx_{\epsilon/10} F$ we conclude that the η satisfies the conclusion of the lemma. $\square$

6. THE MAIN RESULT

Recall that $\mathcal{Z}$ denotes the Jiang-Su algebra.

Theorem 6.1. *If A is an infinite dimensional unital simple AH algebra with no dimension growth (cf. Subsection 2.2), then $A \cong A \otimes \mathcal{Z}$.*

Proof. To prove $\mathcal{Z}$ -stability for A it suffices to prove that for each pair of relatively prime positive integers $p_1, p_2 \geq 2$ there is an approximately central sequence of unital $*$ -homomorphisms $\gamma_n : I_{p_1, p_2} \rightarrow A$ (cf. [13]). In other words, for every $\epsilon > 0$, finite subset F of A , integers p_1, p_2 as above, and finite generating set G for I_{p_1, p_2} consisting of elements of norm at most one, it

will suffice to find a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow A$ such that $\|[\eta(g), f]\| \leq \epsilon$, for all $g \in G$ and $f \in F$.

By assumption, $A = \lim_{i \rightarrow \infty} (A_i, \phi_i)$, where each A_i is semi-homogeneous. There is moreover an $d \geq 0$ such that $\dim(\text{Spec}(A_i)) \leq d$ for every $i \in \mathbb{N}$. Since $\cup_i \phi_{i\infty}(A_i)$ is dense in A , we may assume that F is the image of a finite subset of some A_i ; re-labeling, we simply assume that $F \subset A_1$. Let us observe that A_1 is of the form $pM_N(C(X))p$ with X not necessarily connected.

To prove the theorem, it will suffice to find $j > 1$ and a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow A_j$ such that $\|[\eta(g), \phi_{1,j}(f)]\| \leq \epsilon$ for every $g \in G$ and $f \in F$. We may assume that $\|f\| \leq 1$ for all $f \in F$. Since I_{p_1, p_2} is semiprojective, there is $\epsilon_0 > 0$ smaller than $\epsilon/3$ such that for any completely positive unital map $\mu : I_{p_1, p_2} \rightarrow B$ which satisfies $\|\mu(gh) - \mu(g)\mu(h)\| \leq \epsilon_0$ for all $g, h \in G$ there is a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow B$ with $\|\eta(g) - \mu(g)\| \leq \epsilon/3$ for all $g \in G$. Set $F_0 = F \cup \{p\} \subset M_N(C(X))$.

Set $\gamma = \phi_{1,2} : pM_N(C(X))p = A_1 \rightarrow A_2$. We are going to show that there is a commutative diagram

$$\begin{array}{ccc} pM_N(C(X))p & \xrightarrow{\gamma} & A_2 \\ \downarrow & & \downarrow \\ M_N(C(X)) & \xrightarrow{\gamma_0} & QM_m(A_2)Q \end{array}$$

where γ_0 is a unital $*$ -homomorphism and the vertical arrows are inclusions of full corners. Indeed, if we set $D = M_N(C(X))$ and if $w \in M_m(D)$ is a partial isometry such that $w^*w = \text{diag}(1_D - p, 0, \dots, 0)$ and $ww^* \leq \text{diag}(0, p, \dots, p)$ then $v = \text{diag}(p, 0, \dots, 0) + w \in M_m(D)$ is a partial isometry such that $v^*v = \text{diag}(1_D, 0, \dots, 0)$ and $vv^* \leq \text{diag}(p, p, \dots, p) =: P$. Define $\iota : D \rightarrow PM_m(D)P \cong M_m(pDp)$ by $\iota(a) = vav^*$. Then $\text{id}_m \otimes \gamma : M_m(pDp) \rightarrow M_m(A_2)$ has the property that $(\text{id}_m \otimes \gamma) \circ \iota : D \rightarrow M_m(A_2)$ satisfies $(\text{id}_m \otimes \gamma) \circ \iota(pap) = \text{diag}(\gamma(pap), 0, \dots, 0)$. We set $\gamma_0 = (\text{id}_m \otimes \gamma) \circ \iota : D \rightarrow QM_m(A_2)Q$ where $Q = \gamma_0(1_D)$.

By applying Lemma 5.5 and identifying A_2 and Q with their images in A_j and respectively $M_m(A_j)$ for $j \geq 2$, we find a unital $*$ -homomorphism $\eta_0 : I_{p_1, p_2} \rightarrow QM_m(A_j)Q$ such that $\|[\eta_0(g), \gamma_0(f)]\| \leq \epsilon_0$ for all $g \in G$, and $f \in F \cup \{p\}$. Set $e = \gamma_0(p) = 1_{A_j}$ and observe that the map $e\eta_0(\cdot)e$ is ϵ_0 -multiplicative on G . Therefore by semiprojectivity of I_{p_1, p_2} there is a unital $*$ -homomorphism $\eta : I_{p_1, p_2} \rightarrow eM_m(A_j)e = A_j$ such that $\|\eta(g) - e\eta_0(g)e\| \leq \epsilon/3$ for all $g \in G$. Since $\epsilon_0 < \epsilon/3$ and since F is normalized it follows that

$$\|[\eta(g), \gamma(f)]\| \leq \|[\eta(g) - e\eta_0(g)e, \gamma(f)]\| + \|e\eta_0(g)e, \gamma(f)\| \leq 2\epsilon/3 + \epsilon_0 \leq \epsilon,$$

for all $g \in G$ and $f \in F$. □

Remark 6.2. The no dimension growth hypothesis of Theorem 6.1 can be weakened somewhat. Say that a unital simple AH algebra A has *exponentially slow dimension growth* if for any constant $L > 1$ there is an AH sequence (A_i, ϕ_i) with limit A satisfying

$$\liminf_{j \rightarrow \infty} \max_{1 \leq t \leq n_j} \frac{L^{\dim(X_{j,k})}}{\text{rank}(\mathbf{1}_{A_{j,t}})} = 0.$$

If one replaces the slow dimension growth hypothesis of Lemma 4.2 with the stronger condition of exponentially slow dimension growth, then one can replace the quantity $K \dim(X_{j,k})$ in conclusion (iii) with $K(L')^{\dim(X_{j,k})+2}$ for any constant $L' > 1$. (In the proof, one replaces the numerators equal to $\dim(X_{j,k})$ with $(L')^{\dim(X_{j,k})+2}$.) One can then use exponentially slow dimension growth instead of slow dimension growth in Lemma 5.5—the latter hypothesis is only required for an application of Lemma 4.2. The proof of Theorem 6.1 then goes through as written, with the weakened assumption of exponentially slow dimension growth for A .

There are examples of unital simple AH algebras which have exponentially slow dimension growth but for which one cannot prove bounded dimension growth without the classification theory of AH algebras: the proof of [15, Proposition 5.2] shows that the so-called Villadsen algebras of the first type have exponentially slow dimension growth whenever they have slow dimension growth.

REFERENCES

- [1] Bratteli, O. and Elliott, G. A.: *Small eigenvalue variation and real rank zero*, Pac. J. Math. **175** (1996), 47-59
- [2] Dadarlat, M.: Continuous fields of C^* -algebras over finite dimensional spaces, math.OA/0611405.
- [3] Dadarlat, M., Hirshberg, I., Toms, A. S., and Winter, W.: *The Jiang-Su algebra does not always embed*, arXiv:0712.2020
- [4] Dadarlat, M., Nagy, G., Nemethi, A., and Pasnicu, C.: *Reduction of topological stable rank in inductive limits of C^* -algebras*, Pac. J. Math. **153** (1992), 267-276
- [5] M. Dadarlat, and A. S. Toms: *A Universal Property of the Jiang-Su algebra*, arXiv:0707.4050.
- [6] Elliott, G. A., and Gong, G.: On the classification of C^* -algebras of real rank zero. II. Ann. of Math. (2) **144** (1996), 497-610
- [7] Elliott, G. A., Gong, G. and Li, L.: *On the classification of simple inductive limit C^* -algebras, II: The isomorphism theorem*, Invent. Math. **168** (2007), 249-320
- [8] Elliott, G. A., Gong, G. and Li, L.: *Approximate divisibility of simple inductive limit C^* -algebras*, Operator Algebras and Operator Theory (Shanghai, 1997), 87-97, Contemp. Math. **228**, Amer. Math. Soc., Providence, RI, 1998
- [9] Elliott, G. A., and Toms, A. S.: *Regularity properties in the classification program for separable amenable C^* -algebras*, Bull. Amer. Math. Soc. **45** (2008), 229-245
- [10] Gong, G.: *On the classification of simple inductive limit C^* -algebras I. The reduction theorem.*, Doc. Math. **7** (2002), 255-461
- [11] Goodearl, K. R.: *Riesz decomposition in inductive limit C^* -algebras*, Rocky Mt. J. Math. **24** 1994, 1405-1430
- [12] Jiang, X. and Su, H.: *On a simple unital projectionless C^* -algebra*, Amer. J. Math. **121** (1999), 359-413
- [13] Toms, A. S., and Winter, W.: *Strongly self-absorbing C^* -algebras*, Trans. Amer. Math. Soc. **359** (2007), 3999-4029
- [14] Toms, A. S. and Winter, W.: *$\mathcal{Z}$ -stable ASH algebras*, to appear in Canad. J. Math., arXiv preprint math.OA/0508218 (2005)
- [15] Toms, A. S. and Winter, W.: *The Elliott conjecture for Villadsen algebras of the first type*, arXiv:math/0611059 (2006)
- [16] Winter, W.: *On the classification of simple $\mathcal{Z}$ -stable C^* -algebras with real rank zero and finite decomposition rank*, J. London Math. Soc. (2) **74** (2006), 167-183

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY ST., WEST LAFAYETTE, IN, 47907-2067, U.S.A.

E-mail address: `mdd@math.purdue.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR, 97403, U.S.A.

E-mail address: `ncp@darkwing.uoregon.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, 4700 KEELE ST., TORONTO, ONTARIO, CANADA, M3J 1P3

E-mail address: `atoms@mathstat.yorku.ca`